

SHAPING THE CONTEXT OF HEALTH: A Review of Environmental and Policy Approaches in the Prevention of Chronic Diseases

Ross C. Brownson,¹ Debra Haire-Joshu,²
and Douglas A. Luke³

¹*Department of Community Health and Prevention Research Center,*

²*Department of Community Health and Obesity Prevention Center,*

³*Department of Community Health and Center for Tobacco Policy Research, Saint Louis University School of Public Health, St. Louis, Missouri 63104;*

email: brownson@slu.edu; joshud@slu.edu; dluke@slu.edu

Key Words policy, diet, exercise, physical activity, tobacco

■ **Abstract** Given the growing attention on how environmental and policy interventions can affect chronic disease burden, our objectives are to describe (a) effective and promising interventions to address tobacco use, physical activity, and healthy eating and (b) lessons learned from the literature and practice experience in applying environmental and policy approaches. A total of 17 interventions were reviewed, organized across 3 domains affecting the physical environment/access, economic environment, and communication environment. Many of these interventions are effective. Several lessons are important to consider, such as the need to start with environmental and policy approaches, intervene comprehensively and across multiple levels, make use of economic evaluations, make better use of existing analytic tools, understand the politics and local context, address health disparities, and conduct sound policy research.

INTRODUCTION

The origins of modern public health can be traced back to epidemics of now uncommon infectious diseases such as cholera, plague, and leprosy (127). As these diseases were controlled over the past two centuries, the United States and other countries have experienced a dramatic shift toward chronic diseases as the leading causes of death and disability (16, 141). In 1900, the three leading causes of death were pneumonia and influenza; tuberculosis; and gastritis, enteritis, and colitis. These diseases accounted for nearly one third of all deaths. Today, heart disease, cancer, and stroke are the three leading causes of death, accounting for almost two thirds of all deaths. These and other chronic diseases are characterized

by a complex interaction of risk factors, a noncontagious origin, a long latent period between risk factor exposure and clinical occurrence of disease, a long period of illness, and multiple risk factor etiology (86).

There has been growing attention on how environmental and policy interventions can affect chronic disease burden (18, 48, 70, 85). Such interventions are aimed at changing the physical and sociopolitical environments (111). Environmental and policy approaches are designed to provide opportunities, support, and cues to help people develop healthier behaviors. These approaches serve as an important complement to individual-level programs. For example, environmental and policy changes can benefit all people exposed to the environment rather than focusing on changing the behavior of one person at a time. Alternations in the physical or policy environment may directly affect behaviors (e.g., the influence of the price of tobacco on consumption) or they may alter social norms (e.g., restrictions on smoking in public places). Importantly, environmental and policy approaches are often more permanent than many public health programs focused on individual-level behavioral change.

Although chronic disease practitioners and public health agency leaders value scientific knowledge as a basis for decision making (14), systematic evidence is not always available across a range of risk factors and populations. Practitioners often seek expert advice when searching for effective programs and policies. As an example, an expert panel (i.e., the Task Force on Community Preventive Services) supported by the Centers for Disease Control and Prevention has published *The Guide to Community Preventive Services: What Works to Promote Health?* (i.e., the Community Guide) (145). Evidence-based reviews and summaries (“best practices”) have been useful across many types of chronic disease prevention programs (22, 24).

The primary purposes of this paper are to describe (a) effective and promising interventions to address tobacco use, physical activity, and healthy eating and (b) lessons learned from the literature and practical experience in applying environmental and policy approaches.

A FRAMEWORK FOR THE PREVENTION OF CHRONIC DISEASES THROUGH ENVIRONMENTAL AND POLICY CHANGE

A general conceptual framework (Figure 1) illustrates the relationships between a variety of environmental and policy interventions (those altering the physical environment, the economic environment, or the communication environment); the necessary resources and shifts in norms; changes in risk factors; and long-term chronic disease outcomes. Each of the model domains was derived by common evidence across the target behaviors of tobacco, physical activity, and diet. Although the social environment is not considered as a separate domain, it is directly and indirectly influenced by the domains in Figure 1. For example, the




Figure 1 Conceptual framework for understanding the prevention of chronic diseases through environmental and policy approaches.

presence of healthy foods or places to exercise (in the Physical environment/Access domain) is likely to influence social norms related to healthy eating and physical activity.

In many instances, the outcomes in Figure 1 are not independent. For example, improvements in physical activity and eating patterns would also improve energy balance (28). Although we apply the framework to primary prevention of chronic diseases, this type of model (especially the domains in the rectangles) is likely to be applicable to many other public health issues.

EFFECTIVE AND PROMISING INTERVENTIONS

This review focuses on primary prevention of chronic diseases, i.e., risk reduction among asymptomatic persons to reduce the likelihood of development of chronic disease. We cover three important risk factors: tobacco use, physical inactivity, and unhealthy diet that account for 33% of the annual deaths in the United States (89, 90). There is also an extensive literature on secondary prevention (e.g., cancer screening) and tertiary prevention (e.g., preventing complications and/or

recurrence of cerebrovascular disease). When possible, we rely on the systematic reviews from the Community Guide. The Guide categorizes comparative studies of interventions and takes into account both study design and the quality of the study execution (11, 12, 126). Community Guide recommendations for use are based primarily on the evidence of effectiveness reported in a “best-evidence” subset of the identified studies. As of August 2005, reviews were completed for tobacco and physical activity but are still in progress for diet interventions. When evidence of effectiveness is not clearly documented yet there is high face validity or evidence from another area of public health, the intervention is labeled in this review as a “promising practice.” All interventions are organized within the three domains in Figure 1.

Tobacco Use Reduction

Among the three chronic disease risk factors covered in this review, tobacco has the most robust group of research-tested environmental and policy interventions. It therefore allows us to consider potential mechanisms of change. There are at least three types of environmental effects: policies that shape the (a) behavioral environment, (b) financial environment, and (c) communication environment (Tables 1, 2). Specifically, countries, states, and communities can reduce tobacco harm by reducing opportunities to use tobacco products, making it more expensive to use tobacco (and cheaper to quit using tobacco), and making smoking less attractive, especially to youth.

PHYSICAL ENVIRONMENT/ACCESS

Clean indoor air Clean indoor air laws and regulations reduce exposure to tobacco smoke and restrict opportunities to use tobacco by limiting where and when people can smoke. Since 1971, when the Surgeon General first called for a governmental ban on smoking in public places, a wide variety of national, state, local,

TABLE 1 Framework for understanding environmental and policy approaches to tobacco control

Type of environmental shift	Types of policies	Environmental/Policy mechanism
Physical environment/access	<ul style="list-style-type: none"> ● Clean indoor air ● Restricting youth access 	<ul style="list-style-type: none"> ● Reduce opportunities to obtain and use tobacco products
Economic environment	<ul style="list-style-type: none"> ● Increase cost of tobacco ● Reduce cost of cessation services 	<ul style="list-style-type: none"> ● Make smoking more expensive and quitting less expensive
Communication environment	<ul style="list-style-type: none"> ● Advertising restrictions ● Warning labels 	<ul style="list-style-type: none"> ● Make smoking less attractive

TABLE 2 Summary of environmental and policy interventions to prevent chronic diseases

Category/Intervention	Examples of intervention components	Effectiveness*	Sites/populations involved	Other notes
Tobacco use Clean indoor air	<ul style="list-style-type: none"> • Bans on restrictions on smoking in public spaces and public or private workplaces 	Strong evidence	Wide variety of studies at the organization, community, state, and national level examining the effects of CIA policies in worksites, restaurants, public transportation, schools, and other settings	
Restricting youth access to tobacco	<ul style="list-style-type: none"> • Enactment and enforcement of sales laws directed at retailers • Retailer or community education about youth access • Laws directed at youth purchase, use, or possession of tobacco products 	Insufficient evidence	Large variety of community studies in the United States	Although retail laws and community education can change retail behavior, there is little evidence that these laws and policies can influence youth tobacco use
Taxes and pricing	<ul style="list-style-type: none"> • Increase price of tobacco products through state and federal excise taxes 	Strong evidence	A wide variety of econometric, evaluation and simulation studies examining the effects of price increases. Price effects have been examined in many developed and developing countries	
Reducing cessation costs	<ul style="list-style-type: none"> • Reduce client costs of cessation services 	Sufficient evidence	Small number of studies of insurance coverage. Small number of studies of national health care coverage of cessation services	Studies also found that restrictions on product coverage and availability may inhibit access, even when costs are covered

(Continued)

TABLE 2 (Continued)

Category/Intervention	Examples of intervention components	Effectiveness*	Sites/populations involved	Other notes
Advertising restrictions	<ul style="list-style-type: none"> • Restriction of advertising by media (e.g., television, radio, billboards) • Restriction of marketing to specific populations (i.e., youth) 	Promising evidence	Numerous studies, most often of nationwide bans or restrictions	Data and methodological challenges have limited the ability to detect effects. Only comprehensive bans show noticeable effects
Product warning labels	<ul style="list-style-type: none"> • Requirements to place health warning labels on tobacco products • Requirements to include health warning information in tobacco advertising and marketing 	Promising evidence	Numerous studies of adult and youth in the United States. More recent studies of newer warning labels in other countries	
Physical activity				
Access to facilities	<ul style="list-style-type: none"> • Increased access to fitness centers or community centers • Building walking/biking trails • Focus on recreational activity • Zoning regulations and building codes • Improved street lighting 	Strong evidence	Diverse settings including worksites, universities, federal agencies, and communities	Only one study in ethnically diverse populations
Urban planning & policy	<ul style="list-style-type: none"> • Infrastructure projects that increase the ease and safety of street crossing • Policies to ensure sidewalk continuity 	Sufficient evidence	Includes a variety of geographic scales—from a few blocks to an entire community. Few studies are available among minority populations	Although effect sizes were generally large, all but 3 studies used cross-sectional designs

Transportation policy	<ul style="list-style-type: none"> ● Changing roadway design standards ● Expanding public transportation services ● Subsidizing public transportation 	Insufficient evidence	2 U.S. studies and 1 Canadian study	Other benefits may include air quality, increased commerce, lack of traffic congestion
School-based physical education	<ul style="list-style-type: none"> ● Adding new PE classes ● Lengthening existing PE classes ● Increasing physical activity during PE class without necessarily lengthening class time 	Strong evidence	9 U.S. studies, 1 in Crete, 2 in Australia. 10 studies among elementary school students and 2 studies among high-school students	Primary barrier is the lack of consistent PE policies across states and cities
Economic incentives	<ul style="list-style-type: none"> ● Make funding more available for mixed use projects ● Ensure that gasoline taxes cover all external costs of burning the fuel ● Parking cash-outs where employees are given the cash value of parking ● Performance zoning with the ability to add more footage for pedestrian-oriented streetscapes 	Promising evidence	Some areas have implemented activities in this area, but they have not been systematically evaluated	
Point of decision prompts	<ul style="list-style-type: none"> ● Signs placed by elevators and escalators to motivate people to use nearby stairs ● Messages focus on health benefits or weight loss 	Sufficient evidence	Diverse settings including shopping malls, train/bus stations, and a university library	Some evidence of higher effectiveness in obese individuals

(Continued)

TABLE 2 (Continued)

Category/Intervention	Examples of intervention components	Effectiveness*	Sites/populations involved	Other notes
Healthy eating Food and beverage availability in schools	<ul style="list-style-type: none"> • Food service changes including menu planning, food purchases • Training of food service workers • Improvement in nutrient content, presentation, and quality of school foods • Improved health curriculum • Improve classroom snacks 	Insufficient evidence	Direct comparisons were limited by variations in dietary outcomes including fat, sodium intake, FV consumption, and behavior including food choices and purchases. Evaluation time frames varied from 1 month to 3 years. Interventions included multiple environmental components	Evidence across studies shows consistently positive results, but small effect sizes
Access to healthy ready-to-eat foods	<ul style="list-style-type: none"> • Worksite programs improving ready-to-eat foods purchased in cafeterias or vending machines • Community programs altered ready-to-eat foods provided in restaurants, churches, farmers' markets 	Promising evidence	Multi-component interventions often included health education, labeling or pricing interventions	
Food pricing and incentives	<ul style="list-style-type: none"> • Reduce pricing of healthy vending machine foods • Reduce pricing in cafeterias • Food coupons for FV • Taxation of snack foods 	Promising evidence	Price interventions took place in schools and worksite settings. Additional studies with systematic and robust evaluation designs for pricing interventions are needed	A growing number of studies applying economic principles and research to food behavior and intake show generally consistent and positive results

Nutrition labeling and information	<ul style="list-style-type: none"> ● Point of purchase labeling in cafeterias, worksites, groceries 	Sufficient evidence	Several interventions included reduced pricing and promotion	Labeling of ready-to-eat foods has not been evaluated
Media and marketing policy	<ul style="list-style-type: none"> ● Restaurant menu labeling ● Campaigns promoting education and awareness of healthy FV and foods ● Campaigns limiting advertisements for competitive or foods of minimal nutritional value ● Regulating food advertisements, particularly those targeted to youth 	Insufficient evidence	There is a dearth of studies on dietary campaigns limiting media. Studies have addressed media influence on public education or awareness with mixed results	Media campaigns may be of particular benefit to youth

* When evidence of effectiveness is available from a systematic review (85, 145), it is noted; in other cases, estimates are based on review of the existing literature and the authors' judgment.

and private laws and regulations have been enacted that have dramatically changed the smoking environment in the United States.

The primary effect of clean indoor air regulations is to reduce the exposure of smokers and nonsmokers to harmful secondhand smoke (SS). SS exposure has been linked to a wide variety of harmful health effects that include lung cancer, pneumonia, bronchitis, nasal sinus cancer, and sudden infant death syndrome (93, 128). Smoking bans are, in general, extremely effective in reducing SS exposure. In the studies reviewed by the Community Guide, the median reduction in SS exposure is 60% for self-report measures and 72% for environmental measures (such as nicotine vapor or levels of particulates) (145). In addition to worksite studies, bans on smoking in restaurants and bars also result in lowered exposure rates for community residents (2). Studies have also shown that complete smoking bans are more effective at reducing ETS exposure than smoking restrictions (17).

An important secondary effect of clean indoor air laws is that as the number of smoke-free environments increases, so too does ecological and social pressure to reduce smoking. Cross-sectional analyses have indicated that the prevalence of adult smoking is significantly lower among workers in smoke-free worksites compared with workers at worksites with no smoking restrictions (19). Stronger evidence for the effects of clean indoor air policies on smoking behavior is found in a set of ten experimental or prospective studies reviewed in the Community Guide (145). These studies all show that implementation of strong clean indoor air regulations results in a reduction in rates of cigarette consumption, smoking prevalence, and increased cessation.

Restricting Youth Access to Tobacco Although it is illegal to sell tobacco products to minors, the vast majority (82%) of adults in the United States who have ever smoked tried their first cigarette before age 18, and 53% became daily smokers before age 18 (129). The purpose of youth access policies is to reduce the sales of tobacco products to teenagers and thus lower smoking prevalence by preventing or delaying smoking initiation.

Although states with more extensive youth access policies have significantly lower youth smoking rates (81), it has been difficult to demonstrate a causal link between implementation and enforcement of youth access policies and subsequent reductions in youth smoking. Programs that implement community and retailer education, sales monitoring, and policy enforcement can achieve significant changes in retail sales behavior, but this does not necessarily translate into changes in smoking behavior. Recent reviews of youth access interventions and purchase, use, and possession policies have summarized the lack of strong evidence, and they have argued that the difficulty in program monitoring, the ability of teens to acquire cigarettes from social sources, and the high cost of policy enforcement all suggest that tobacco control advocates should pursue other policies that are more cost-effective (132).

ECONOMIC ENVIRONMENT

Taxes and pricing One of the simplest and most effective ways to reduce smoking and tobacco use is to raise the price of tobacco products. Economic behavioral theory suggests that if cigarettes cost more, then fewer people will start smoking, existing smokers will smoke less often, and more smokers will attempt to quit. The primary way to influence price is through federal and state excise taxes. In 2005, the federal tax rate was 39 cents per pack and the median state tax rate was just under 70 cents per pack (4).

A large number of econometric studies have examined the relationship between tobacco prices and tobacco use (called elasticity) for adults and youth in both developed and developing countries. For adults, a 10% increase in cigarette prices is estimated to produce a 3% to 5% reduction in demand (77, 135). There is an emerging consensus that youth and young adults are even more sensitive to price than are adults (77). Estimates range from around 7% to 8% (51, 121) to as high as 14% (78). Recent studies have shown that higher prices affect smoking by lowering prevalence rates, lowering quantity of cigarettes smoked, decreasing the duration of smoking, and reducing initiation (30, 77). Finally, international studies have shown that in developed countries price elasticity is similar to that found for the United States, whereas developing and low-income countries show greater price sensitivity (60, 77).

In addition to their proven effectiveness, price increases through tax policy are attractive because of their broad reach. National and state excise taxes affect every tobacco consumer. Some have worried that increased prices would result in tax evasion and avoidance through increased purchase of black market cigarettes, increased use of the Internet to purchase low-priced cigarettes, or increased cross-border cigarette purchasing and smuggling. However, although studies have shown that consumers will purchase lower-priced cigarettes when they are readily available (e.g., if they live very close to an Indian reservation or bordering state with lower prices) (62), most studies show that, in fact, tobacco consumers find it difficult to avoid higher cigarette prices when they are introduced (112). For example, only around 5% of California smokers avoided higher prices by changing their purchase habits after a 50 cent/pack tax increase (36).

Reducing cessation costs Another way to shape the financial climate for tobacco control is to reduce the costs for cessation products and services for established smokers. Out-of-pocket costs for cessation are reduced primarily by increased public or private insurance coverage. A small number of studies have shown that reducing cessation costs can increase the number of people who use cessation services by approximately 7%, and increase the number of people who successfully quit using tobacco by approximately 8% (145).

Although insurance coverage of cessation services has been shown to be cost-effective, these services are still not widely covered in United States insurance and managed care systems (77). In the United Kingdom, the National Health System

has started to cover cessation pharmacotherapies, and early evidence suggests that this coverage has increased use of services, quit attempts, and short-term quit rates (77).

COMMUNICATION ENVIRONMENT

Advertising restrictions Tobacco products are one of the most heavily advertised and promoted products on the planet. Bans or restrictions on tobacco advertising have played an important role in tobacco control policy. In 1970 tobacco advertising was banned on television and radio. Billboard advertising of tobacco products was banned in 1999 as part of the Master Settlement Agreement, and as part of the same agreement tobacco companies agreed to stop advertising their products directly to youth.

Despite these advertising restrictions, there has been little evidence that partial restrictions have any effect on youth or adult smoking. One challenge is methodological—it is difficult to obtain relevant data, and the inability to examine large variations in tobacco advertising may make it impossible to detect causal relationships (77). Partial bans also leave open other marketing opportunities for the tobacco industry. For example, there is evidence that after the Master Settlement Agreement tobacco companies increased their direct marketing and bar promotion activities in an attempt to continue to reach young smokers (71).

However, recent data suggest that countries that adopt comprehensive advertising bans can expect to reduce tobacco use by approximately 5.4% (107). The societal cost to implement such national comprehensive bans is quite low, although there is stiff political pressure against adopting such bans.

Product warning labels The United States has required health warning labels on cigarettes since 1965, and most other countries require some type of labeling that supplies product or health information. Studies of warning labels like those used in the United States show that they have little effect. Warning labels are easily ignored, hard to recall, and do not stand out from other packaging elements (72). However, recent evidence from countries such as Canada and Australia, which have required larger, more colorful and graphic warning labels, indicates that labels can influence attitudes and behavior (7). For example, in Canada 44% of smokers were more motivated to quit after seeing the labels, and 21% of smokers who were tempted to have a cigarette did not because of the warning labels (83).

Promotion of Physical Activity

Specific strategies to promote physical activity with environmental and policy approaches were first proposed approximately a decade ago (70). At that point, the empirical basis for these interventions was quite limited. Since then, a substantial body of correlates research (61, 98, 106), along with a smaller number of intervention studies, has been published. Examples of environmental and policy

approaches to increase physical activity include walking and bicycle trails, funding for public facilities, zoning and land use that facilitates activity in neighborhoods, building construction that encourages activity, policies and incentives promoting physical activity during the workday, and policies requiring school-based physical education (66, 70, 109).

This section describes approaches to promotion of physical activity in six specific categories that bridge the domains in Figure 1. For some intervention categories there is a substantial and consistent body of literature, whereas for others the literature is sparse (although sometimes promising).

PHYSICAL ENVIRONMENT/ACCESS

Access to facilities Strategies often include providing access to facilities that are not currently available to the local population. A review of 19 studies in the physical activity and health literature showed consistent associations of accessibility of recreational facilities, opportunities to be active, and aesthetic qualities with physical activity in adults (61).

Interventions reviewed in the Community Guide involved the efforts of work-sites, coalitions, agencies, and communities to create or provide access to places and facilities where people can be physically active. For example, interventions include providing access to nearby fitness centers or to weight and aerobic fitness equipment in community centers and creating walking trails. Many of these studies also incorporated components such as training on equipment, health behavior education and techniques, seminars, counseling, risk screening, health forums and workshops, referrals to physicians or additional services, health and fitness programs, and support or buddy systems.

Quantitative estimates of effect are based on ten studies (66). In eight arms from five studies the median net change in aerobic capacity was 5%. In four arms from three studies the median net change in the frequency of physical activity was 48%.

Urban planning and policy There have been dramatic changes in the urban landscape over the past 50 years (15). Muller has described the period from 1945 to the present as the "Freeway Era" in which the automobile is no longer a luxury but has become essential for commuting, shopping, and socializing (92). This trend also contributed to the advent of the suburban ring and to the accompanying freeway segments that now girdle most central cities in the United States. The migration to suburban environments is closely linked with the evolution of zoning policies over the past century (110). For example, landmark cases such as *Euclid v. Amber Realty* (1926) established the importance of local zoning laws in shaping the patterns of growth in urban areas.

Researchers in transportation and urban planning have examined the relationship between community design variables and walking or cycling for transportation. A review of 14 studies consistently shows that people walk and cycle more when their neighborhoods have higher residential density, a mixture of land uses

(e.g., shops are within walking distance of homes), and connected streets (e.g., grid-like pattern instead of many cul-de-sacs) (106).

The Community Guide identified two types of interventions through urban planning and policy (57). The first set are urban design strategies that involve street-scale changes and the second are land use policies that support physical activity in small geographic areas, generally limited to a few blocks. Another closely related group of interventions focus on community-scale changes that are similar to street-scale changes but involve a much larger geographic area (e.g., an entire community).

Among the total group of 19 studies, 16 were cross-sectional and 3 had pre- and post-designs. The metrics used varied widely, including outcomes such as percent of the population walking to work; percent of household pedestrian trips compared to transit and auto trips; average frequency of walking trips to shopping area each month; and self-reported number of minutes walked per week or per month, percent of trips made by walking or biking, and distance and duration of walking trips. Median net effects ranged from 35% for street-scale interventions to 161% for community-scale interventions.

Transportation policy Transportation policy can have substantial impacts on travel choice (e.g., walking versus driving). Changes in transportation policy may influence the rate of physical activity and may also be beneficial to air quality and traffic congestion.

This group of interventions rely on policy measures such as roadway design standards, expansion of public transportation services, or subsidization of public transportation (i.e., transit passes). In the review of the Community Guide only three studies were identified for this intervention strategy. Due to too few qualified studies, this intervention strategy has insufficient evidence to make any type of recommendation. This is an important area for future research.

School-based physical education Interventions in this category involve modified curricula and policies to increase the amount of time students spend in moderate or vigorous activity while in physical education (PE) classes. This modification is accomplished in numerous ways including (a) adding new (or additional) PE classes, (b) lengthening existing PE classes, or (c) increasing moderate to vigorous physical activity of students during PE class without necessarily lengthening class time. Examples of specific interventions include changing the activities taught (e.g., substituting soccer for softball) or modifying the rules of the game so that students are more active (e.g., having the entire team run the bases together if the batter makes a hit). Many of these interventions also included the presentation of information on cardiovascular disease prevention, rendering it difficult to separate the effects of health education and modified PE.

Among the 13 studies reviewed, outcome measures included energy expenditure, percent of class time spent in activity, minutes spent in activity, and self-reported type and frequency of physical activities outside of school (66). Five

arms from four studies showed increases in the amount and percent of time spent in physical activity in PE classes. The net increase in the amount of PE class time spent in physical activity was 50%. The net increase in the percent of class time in physical activity was 10%, with an additional study reporting a 762% increase from a very small baseline value. Three arms from two studies showed increases in energy expenditure as well. Fourteen arms from eleven studies showed increases in aerobic capacity, with a median of 8%. There is also evidence that these school-based changes can be maintained and institutionalized after research funding ends (58).

ECONOMIC ENVIRONMENT

Economic incentives There is growing interest in understanding the economic impact of physical inactivity and the economic interventions that might be effective in promoting activity. The economic consequences of inactivity are substantial, with direct medical expenses estimated at \$76 billion in year 2000 in the United States (102).

When considering economic interventions, it is important to consider that economics, especially those influencing the transportation infrastructure, strongly influence individual-level transportation choices (87). A comprehensive set of potential economic intervention strategies was published by Pratt and colleagues (101). These interventions are closely related to those in transportation policy noted above; however, they specifically focus on how financial incentives and disincentives influence physical activity patterns.

There are few empirical data showing the effectiveness and magnitude of effects of these economic interventions. At this point, they represent a promising set of strategies that should be implemented and evaluated (so-called natural experiments).

COMMUNICATION ENVIRONMENT

Point-of-decision prompts Cues to make use of an environmental feature are important in promoting physical activity. The use of cues comes via point-of-decision prompts, e.g., signs placed by elevators and escalators to motivate people to use nearby stairs. Messages on the signs recommend stair use for health benefits or weight loss. Signs are thought to be effective in one of two ways: by reminding people already predisposed to becoming more active, for health or other reasons, about an opportunity at hand to be more active or by informing them of a health benefit from taking the stairs.

The Community Guide reported on six studies in its 2002 review. These studies were conducted between 1985 and 2000. Across this group of studies, baseline rates of stair use were generally low, with all but one under 12% (range: 4.8% to 39.6%). The range of effect sizes varied from a 5.5% net increase to 128.6%. Since the publication of Kahn et al. (66), there are at least ten additional original studies in the peer-reviewed literature. These newer studies begin to address effects across

ethnic groups (25) and the effects of poster size and content in promotion of stair use (69).

Promotion of Healthy Eating

Trends in diet quality show increases in consumption of energy-dense, low-nutrient foods, associated with the development of a variety of chronic diseases (43, 65, 96, 139). To reverse these trends, public health practitioners have begun to focus more on changing environmental and policy factors that promote healthy eating (8, 53). These interventions fall within five categories: pricing or economic approaches, nutrition labeling or information strategies, availability of nutritious foods in schools, access to healthier ready-to-eat foods, and media or public education approaches. These are organized according to the broad headings in Figure 1.

PHYSICAL ENVIRONMENT/ACCESS

Food and beverage availability in schools Policies to assure a healthy school environment can have repercussions on the approximately 52 million students who attend school and the 28 million children participating in the National School Lunch Program (NSLP) (26). Competitive foods and beverages offered à la carte in schools meet minimal standards for nutrition content. These à la carte foods are easily accessible and affordable to children in 43% of elementary schools, 89% of middle schools, and 97% of high schools that have vending machines or other school food services outside of school lunch programs (137).

Much of the research in schools is designed to improve availability of healthy foods and beverages by combining multicomponent behavioral and environmental interventions. These approaches generally improve nutrition intake by altering food services, preparation and choices in cafeterias and vending machines; offering promotional activities for healthy foods and beverages; enhancing nutrition content of classroom curricula; and informing and educating parents and teachers (35, 55, 56, 64, 82, 95, 100, 125). Policies in accordance with CDC recommendations for school-based healthy eating programs have also been shown to be effective in improving health outcomes (131). School-based interventions report small but consistent improvements in student intake of fruits and vegetables (FV) (52, 82, 95, 100), decreases in fat intake (35, 55, 125), and choices of healthier food options (59).

A report by the Institute of Medicine in 2005 concluded that availability of healthy foods alone may not be sufficient to promote positive choices for children when competitive foods high in fat and sugar are easily accessible (26). Environmental interventions are most effective that make healthy options available while also restricting the availability of competitive foods (26). The inclusion of pricing strategies and assuring information about healthy foods also represents promising environmental strategies for improving intake of youth in school settings (119).

Access to healthy ready-to-eat foods Foods obtained away from home are often ready to eat and prepared by another vendor, meaning that the consumer has little control over nutrient content or portion size (41). Ready-to-eat foods typically offer portion sizes 2 to 5 times larger than those found in previous decades and account for over one third of the total energy intake of the population (76, 134, 144). The convenience of ready-to-eat foods in community restaurants, fast food venues, or local stores, as well as in worksite cafeterias and vending outlets, plays a role in the population trend of increasing caloric intake.

Environmental intervention studies addressing ready-to-eat foods have sought to improve the quality of offerings across a variety of settings. Several worksite interventions have reported improvements in food preparation and access to healthy ready-to-eat foods as effective strategies for increasing FV consumption or reducing fat intake (6, 99, 115). Community-based studies have evaluated improved FV or dietary composition in church-based settings (20), restaurants (27, 32, 50), and hotels or entertainment venues (138). These studies varied in research design, length to follow-up (from 1 month to 4 years) and primary outcomes. However, in general, each reported outcomes of improved food preparation or access; three of the studies increased customer requests for healthy entrees (27, 32, 50). One case study reported strategies that resulted in the altered policies that assured neighborhood access to healthy food retailers (136). To date, there is a dearth of environmental studies testing interventions in fast food restaurant venues.

ECONOMIC ENVIRONMENT

Food pricing and incentives Reviews of agricultural and nutritional food data and policies note increases in food intake occurred in conjunction with the commercialization of the food supply. This is defined by incentives to improve sales through marketing and increasing demand for food (123, 124). Between 1985 and 2000 the price of fresh fruits and vegetables increased 118% compared with a 20% to 46% increase in the price of calorie-dense foods and beverages of minimal nutritional value (e.g., snacks and soda) (31). Economic principles have established that decreases in food pricing increase consumption, as evidenced by supersizing, which may partially account for the reported increase in intake of energy-dense foods (38, 88, 143).

Recent intervention research has applied the principle of price reduction to sales and consumption of low-fat snacks in vending machines. A series of studies in schools and communities demonstrated that progressive lowering of prices of low-fat snacks yielded significant increases in sales coinciding with the amount of price reduction (45). A 50% price reduction of fresh fruits and vegetables in school cafeterias resulted in two- to fourfold increases in sales of carrots and fresh fruit (54). A review by Glanz and colleagues reported positive impacts from price reduction coupons used by seniors and WIC participants at farmers' markets (49). Others have also reported enhanced impact on dietary intake when coupons were combined with education (5).

Increases in food or beverages taxes have been suggested as a means of decreasing consumption and increasing revenue for health promotion programs. Studies examining the expected impact of snack food taxation concluded a 1% tax would only minimally affect consumption of salty snacks (e.g., potato chips) by 0.28 ounces or 42 calories per person per year. A 20% tax would reduce consumption by 5.54 ounces per person per year, or 830 calories, and would generate \$40 to \$100 million in tax revenues (73, 74). As of 2000, 19 states and cities have implemented policies on taxing snack foods, soft drinks, candy or chewing gum; however, impact data on these policy initiatives are not yet available (13).

Environmental interventions targeting food costs as a means of altering intake are promising, but the lack of robust research studies to date on outcomes of these policies limits our ability to predict their effectiveness.

COMMUNICATION ENVIRONMENT

Nutrition labeling and information Empirically sound information is critical if the public is to make informed choices regarding dietary intake. The Nutrition Labeling and Education Act was implemented in 1990 (45) to provide the public with accurate information and food facts to guide choices. Labels are available on over 95% of all processed foods and influence purchasing patterns of the public (94). However, nutrition labels are frequently lacking on foods eaten away from home, at restaurants, vending machines, or fast food venues, which account for 46% of all food dollars spent (31, 33, 79, 103). This suggests that consumers are not fully informed about the content of much of what they are eating.

Reviews have been conducted of worksite, community, and school interventions addressing the effectiveness of nutrition labeling on purchase or intake (8, 47, 52, 85). Labeling of vending machine foods has been shown to further enhance consumption of reduced price low-fat foods in school settings (42, 44). A review by Matson-Koffman et al. identified 11 studies that included labeling in interventions that showed improvement in the selection of healthy foods in worksites and community settings that included restaurants and grocery stores (85).

Glanz and colleagues (47, 49) reviewed restaurant-based environmental and policy interventions including point of purchase (POP) studies defined by the use of shelf labels or signage to specify healthy food choices (47, 49, 65). Although several POP studies lacked rigorous design elements, the results generally supported feasibility and improved knowledge. Other studies of more rigorous design reported supermarket POP programs significantly reduced intake of fat in meats and fried foods (27), increased sales of labeled grocery foods (49), and improved knowledge and intake of target foods (85).

There is limited information to validate the effect of labeling on the nutrient intake of the population. However, knowledge and behavioral outcomes associated with the current nutrition-labeling program suggest merit in policies that extend labeling to include foods eaten away from home.

Media and marketing policies Advertising and marketing are intrinsically linked to the food supply and intake of the population (46). Food-marketing practices have emphasized the convenience, low cost, and increased availability of energy-dense foods (65). The food industry spends \$50 per person per year to advertise foods compared with \$1.50 spent by the USDA (46). Food and beverage advertisers collectively spend 10 to 12 billion dollars a year to reach children and youth who see 40,000 TV ads per year promoting snack foods and beverages (26, 67). This repeated exposure to food commercials influences children's preferences and food requests, while ads can also contribute to confusion about the relative health benefits of certain foods.

Media campaigns designed to balance or enhance health messages addressing dietary change have often been conducted in combination with other multicomponent interventions (9, 29, 32). Reviews of several studies targeting nutrition showed that results have been mixed, with improvements cited in the number of adults who switched to low-fat milk (104), or achieved small improvements in FV (39, 117), or in fat consumption (40). Others found improvements in awareness or intention to alter diet but not in consumption (130). Campaign effectiveness is influenced by the intensity of message exposure, reach, timing, and cost.

There is limited evidence of effectiveness for nutrition policies that regulate media messages addressing unhealthy food intake. However, given the success of this approach in other areas (e.g., tobacco control), regulation of media messages remains a promising strategy and an important area for future research.

CHALLENGES AND LESSONS LEARNED

Chronic diseases, including cardiovascular diseases, cancers, diabetes, stroke, and respiratory diseases, account for the majority of the global disease burden (140). Other reviews (85, 145) and ours suggest there are numerous approaches with significant potential for reducing the incidence of chronic diseases.

In developing and implementing these interventions, public health agencies and other community-based organizations face several particular challenges. First, chronic diseases are often not seen as a crisis compared with other health threats such as infectious diseases. Second, the "pay-off" for chronic disease prevention efforts occurs in future years not in weeks or months. Third, the public often shows more concern about involuntary risks (e.g., potential exposure to a chemical waste site) than about voluntary risks (e.g., cigarette smoking), even though the voluntary risks account for the largest burden from chronic diseases. Fourth, many communities lack the local data on chronic diseases and their risk factors for priority setting and program evaluation. This issue is beginning to be addressed (e.g., statewide risk factor data) but remains a serious constraint at the county, city, and neighborhood levels. And finally, sufficient resources have not been committed to chronic disease control efforts. Public health funding dedicated to chronic

disease activities is disproportionately low in relation to the public health burden of chronic diseases (21).

In spite of these challenges, there have been numerous successes in implementing environmental and policy interventions to reduce chronic disease risk (22–24, 142). From these experiences, we can draw useful lessons.

The issues that follow are common across all three risk factors. There also are certain differences that deserve mention. The “rules of engagement” for the industries involved need to be carefully considered (142). It is unlikely that constructive dialogue with the tobacco industry will occur, yet for other areas (e.g., physical activity), well-defined partnerships with industry [e.g., athletic shoe manufacturers (97)] may prove beneficial. There are also differences in risk behaviors (e.g., smoking) that should be totally avoided versus necessary behaviors (e.g., eating) that represent a range of risk. Finally, there is a much larger body of evidence on intervention-related studies for tobacco than for the other two behaviors.

Start with Environmental and Policy Interventions

Based on public health movements, policy and environmental change is key to initiating and sustaining systematic change (34). In many cases, control of chronic diseases is most effective if environmental and policy approaches are the earliest focus of change. These approaches can be low cost, high reach, and tend to provide supportive environment for later targeted interventions. For example, if one is implementing a community-based program to promote energy balance (healthy eating and physical activity), there is little chance of effectiveness if there is a lack of access to healthy foods or places for activity. This fits well with the World Health Organization’s statement in its Ottawa charter that “Healthy choices need to be the easy choices” (142). We suggest that an opportune starting point for environmental and policy interventions should involve the three domains in our framework (the left-hand rectangles in Figure 1).

Think Comprehensively and Across Multiple Levels

Drawing on the process of assembling systematic reviews, as well as evidence from state and local chronic disease-prevention efforts (11, 22, 80, 114, 145), comprehensive interventions that address multiple levels of an ecological framework are more effective. For example, to address childhood obesity (through healthy eating and increased activity), multilevel approaches are needed to include schools that sell or serve only healthy foods and beverages, stronger policies for school physical education, and zoning and subdivision ordinances that support and encourage walking and bicycling (26).

Make Use of Economic Evaluations

There is a considerable body of information on economic evaluation of tobacco control interventions (60) and growing evidence for physical activity and diet.

In tobacco control, the estimated net economic benefit of a national ban on smoking in nonresidential buildings was between \$42 and \$78 billion per year (91). In one of the few economic evaluations in physical activity, Wang and colleagues (133) examined the cost of trail development per trail user in Lincoln, Nebraska. The annual cost per user was \$235 (range = from \$83 to \$592), whereas per capita annual medical cost of inactivity was \$622. Studies like these supply powerful information for health advocates and policy makers.

Make Better Use of Existing Tools

This review highlights the importance of systematic reviews, such as those found in the Community Guide. Another important tool is Cancer Control PLANET (1). The PLANET portal provides access to data and resources that can help planners, program staff, and researchers to design, implement and evaluate evidence-based cancer control programs. Active training and dissemination efforts are needed to increase the use of tools like the Community Guide and PLANET, which are currently underutilized (14, 68, 84).

Understand Local Context

When implementing environmental and policy interventions, it is important to keep in mind that intervention effectiveness does not necessarily equate with intervention feasibility (e.g., a highly effective intervention may be very expensive to implement). To better understand feasibility, assessment of local context is essential. The local context for an intervention should be assessed in conjunction with local data (37, 122) and with information on how to apply interventions found in systematic reviews like the Community Guide (145).

Understand the Politics

Policy interventions are by nature political. Policy makers have to sell, argue, advocate, and get re-elected in light of available political capital. Their interests often tend to be shorter-term and are keyed to an election cycle. Even in the presence of sound scientific data on how to prevent chronic diseases, ideas are sometimes not ready for policy action—owing to lack of public support or competing policies. Practitioners working in governmental agencies have to learn how to influence policy development without direct lobbying. This often involves working in coalitions that can mobilize support and advocate for specific issues. These broad-based coalitions have been essential in achieving progress in tobacco control (142).

Build New and Nontraditional Partnerships

The importance of transdisciplinary partnerships has long been recognized in public health (63, 105). Often the goal of those partnerships is to enhance the work of the public health partners. In many cases, successful implementation of

environmental and policy interventions will require new skills and nontraditional partnerships with people and organizations not working directly in public health. For example, to address the major structural barriers to physical activity in U.S. cities, urban planners, transportation experts, and persons working in parks and recreation are essential collaborators in developing the environment and the political will for activity-friendly communities (108). To address the food environment, partnerships might be explored with organizations engaged in food packaging and processing, marketing, or sales.

Address Health Disparities

In both developed and developing countries, poverty is strongly correlated with poor health outcomes (120). Most of the existing intervention literature has been conducted among ethnic majority populations and higher-income populations. A challenge for the application of environmental and policy strategies involves a better understanding of how interventions apply within populations with large health disparities. Policy interventions have the potential to equalize the environment in a way that may significantly reduce the growing disparities.

Learn from Other Parts of the World

There are many examples of successful environmental and policy approaches worldwide that can provide guidance in the United States. For example, European studies have suggested the importance of policy orientation (i.e., a collection of strategies, policy statements, and committee reports that express the will of the government). Countries that place a higher emphasis on opportunities and infrastructure for physical activity appear to have higher rates of activity (118). Policies designed to inform and alter public purchasing patterns (i.e., adoption of labeling systems countrywide, information on soft drink versus water campaigns) have improved health outcomes of the population (9).

Conduct More and Better Policy Research

We still have a great deal to learn about environmental and policy approaches. Rich opportunities for policy research may take a number of forms: (a) identifying relevant policies (surveillance); (b) understanding the determinants of establishing policy; (c) exploring the process of developing and establishing policy; and (d) assessing the outcomes of policy implementation. In these studies, the policy can be either the independent or dependent variable.

In mainstream epidemiology, the most rigorous design for hypothesis testing is the randomized controlled trial (75). However, a randomized design is seldom useful in policy research because the scientist cannot randomly assign exposure (the policy). Therefore, quasi-experimental designs (e.g., ecologic studies, time-series designs) are likely to be more useful for many policy-relevant issues. Policy research can still be sophisticated in the absence of randomized designs (e.g., the

multilevel nature of policy—federal, state, local—leads to multilevel modeling strategies) (3, 10, 113, 116).

CONCLUSION

There is now a considerable body of evidence showing that environmental and policy interventions are effective in preventing chronic disease risk factors. The evidence is strongest for tobacco use reduction, yet sizeable and growing in volume for promotion of physical activity and healthy eating. The large and increasing burden of chronic diseases can be reduced through systematic and sustained implementation of environmental and policy interventions. Before implementing an array of individual-level programs to prevent chronic diseases, practitioners should consider the power of environmental and policy approaches to set the stage for other interventions.

ACKNOWLEDGMENTS

This project was partially funded through the Centers for Disease Control and Prevention award U48 DP00060-01 (Health Promotion and Disease Prevention Research Center). The authors appreciate the helpful feedback from Drs. Laurie Anderson and Jonathan Fielding.

**The Annual Review of Public Health is online at
<http://publhealth.annualreviews.org>**

LITERATURE CITED

1. 2005. Cancer Control PLANET. *Links to comprehensive cancer control resources for public health professionals*. <http://cancercontrolplanet.cancer.gov/>
2. Albers AB, Siegel M, Cheng DM, Rigotti NA, Biener L. 2004. Effects of restaurant and bar smoking regulations on exposure to environmental tobacco smoke among Massachusetts adults. *Am. J. Public Health* 94:1959–64
3. Alexander LN, Seward JF, Santibanez TA, Pallansch MA, Kew OM, et al. 2004. Vaccine policy changes and epidemiology of poliomyelitis in the United States. *JAMA* 292:1696–701
4. Am. Lung Assoc. 2005. *State legislated actions on tobacco control*. <http://slati.lungusa.org/>
5. Anderson J, Bybee D, Brown R, McLean D, Garcia E, et al. 2001. 5 a day fruit and vegetable intervention improves consumption in a low income population. *J. Am. Diet. Assoc.* 101:195–202
6. Beresford SA, Thompson B, Feng Z, Christianson A, McLerran DF, Patrick DL. 2001. Seattle 5 a day worksite program to increase fruit and vegetable consumption. *Prev. Med.* 32:230–38
7. Borland R. 1997. Tobacco health warnings and smoking-related cognitions and behaviours. *Addiction* 92:1427–35
8. Bowen DJ, Beresford SA. 2002. Dietary interventions to prevent disease. *Annu. Rev. Public Health* 23:255–86
9. Brannstrom I, Weinehall L, Persson L, Wester P, Wall S. 1993. Changing social

- patterns of risk factors for cardiovascular disease in a Swedish community intervention programme. *Int. J. Epidemiol.* 22:1026–37
10. Brewer G, deLeon P. 1983. *The Foundations of Policy Analysis*. Homewood, IL: Dorsey
 11. Briss PA, Brownson RC, Fielding JE, Zaza S. 2004. Developing and using the Guide to Community Preventive Services: lessons learned about evidence-based public health. *Annu. Rev. Public Health* 25:281–302
 12. Briss PA, Zaza S, Pappaioanou M, Fielding J, Wright-De Agüero L, et al. 2000. Developing an evidence-based Guide to Community Preventive Services—methods. The Task Force on Community Preventive Services. *Am. J. Prev. Med.* 18:35–43
 13. Brownell K, Jacobsen M. 2000. Small taxes on soft drinks and snack foods to promote health. *Am. J. Public Health* 90: 854–57
 14. Brownson RC, Ballew P, Dieffenderfer B, Haire-Joshu D, Heath GW, et al. 2006. What contributes to dissemination by state health departments of evidence-based interventions to promote physical activity? *Am. J. Prev. Med.* In press
 15. Brownson RC, Boehmer TK, Luke DA. 2005. Declining rates of physical activity in the United States: What are the contributors? *Annu. Rev. Public Health* 26:421–43
 16. Brownson RC, Bright FS. 2004. Chronic disease control in public health practice: looking back and moving forward. *Public Health Rep.* 119:230–38
 17. Brownson RC, Hopkins DP, Wakefield MA. 2002. Effects of smoking restrictions in the workplace. *Annu. Rev. Public Health* 23:333–48
 18. Brownson RC, Koffman DM, Novotny TE, Hughes RG, Eriksen MP. 1995. Environmental and policy interventions to control tobacco use and prevent cardiovascular disease. *Health Educ. Q.* 22:478–98
 19. Burns DM, Shanks TG, Major JM, Gower KB, Shopland DR. 2000. Restrictions on smoking in the workplace. In *NCI Population Based Smoking Cessation*, pp. 99–128. *Proc. Conf. What Works to Influence Cessat. Gen. Popul.* NCI Monogr. 12. Bethesda, MD: US DHHS
 20. Campbell MK, Demark-Wahnefried W, Symons M, Kalsbeek WD, Dodds J, et al. 1999. Fruit and vegetable consumption and prevention of cancer: the Black Churches United for Better Health project. *Am. J. Public Health* 89:1390–96
 21. Cent. Dis. Control. 1997. Resources and priorities for chronic disease prevention and control, 1994. *MMWR* 46:286–87
 22. Cent. Dis. Control Prev. 1999. *Best Practices for Comprehensive Tobacco Control Programs*. Atlanta, GA: Natl. Cent. Chronic Dis. Prev. Health Promot., Off. Smok. Health
 23. Cent. Dis. Control Prev. 1999. Ten great public health achievements—United States, 1900–1999. *MMWR* 48:241–43
 24. Cent. Dis. Control Prev. 2003. *Promising Practices in Chronic Disease Prevention and Control. A Public Health Framework for Action*, Atlanta, GA: Cent. Dis. Control Prev.
 25. Coleman KJ, Gonzalez EC. 2001. Promoting stair use in a US-Mexico border community. *Am. J. Public Health* 91: 2007–9
 26. Comm. Prev. Obesity in Children and Youth. 2004. *Preventing Childhood Obesity. Health in the Balance*. Washington, DC: Natl. Acad.
 27. Croft J, Temple S, Lankenau B, Heath G, Macera C, et al. 1994. Community intervention and trends in dietary fat consumption among black and white adults. *J. Am. Diet. Assoc.* 94:1284–90
 28. Dietz WH, Gortmaker SL. 2001. Preventing obesity in children and adolescents. *Annu. Rev. Public Health* 22:337–53
 29. Dixon H, Borland R, Segan C, Stafford H,

- Sindal C. 1998. Public reaction to Victoria's "2 fruit 'n' 5 veg every day" campaign and reported consumption of fruit and vegetables. *Prev. Med.* 27:572-82
30. Douglas S. 1998. The duration of the smoking habit. *Econ. Inq.* 36:49-64
31. Drewnoski A, Darmon N. 2005. Food choices and diet costs: an economic analysis. *Am. Soc. Nutr. Sci.* 135:900-4
32. Dunt D, Day N, Pirkis J. 1999. Evaluation of a community-based health promotion program supporting public policy initiatives for a healthy diet. *Health Promot. Int.* 14:317-27
33. Ebbeling C, Sinclair K, Pereira M, Garcia-Lago E, Feldman H, Ludwig D. 2004. Compensation for energy intake from fast food among overweight and lean adolescents. *JAMA* 291:2828-33
34. Economos CD, Brownson RC, DeAngelis MA, Novelli P, Foerster SB, et al. 2001. What lessons have been learned from other attempts to guide social change? *Nutr. Rev.* 59:S40-56; discussion S7-65
35. Ellison R, Goldberg R, Witschi J, Capper A, Puleo E, Stare F. 1990. Use of fat-modified food products to change dietary fat intake of young people. *Am. J. Public Health* 80:1374-76
36. Emery S, White MM, Gilpin EA, Pierce JP. 2002. Was there significant tax evasion after the 1999 50 cent per pack cigarette tax increase in California? *Tob. Control* 11:130-34
37. Fielding JE, Frieden TR. 2004. Local knowledge to enable local action. *Am. J. Prev. Med.* 27:183-84
38. Finkelstein E, Ruhm C, Kosa K. 2005. Economic causes and consequences of obesity. *Annu. Rev. Public Health* 26:239-57
39. Foerster S, Kizer K, DiSogra L. 1995. California's "5 a day-for better health" campaign: an innovative population-based effort to effect large-scale dietary change. *Am. J. Prev. Med.* 11:124-31
40. Fortmann S, Flora J, Winkleby M, Schooler C, Taylor C, Farquhar J. 1995. Community intervention trials: reflections on the Stanford Five-City Project Experience. *Am. J. Epidemiol.* 142:576-86
41. French S. 2005. Public health strategies for dietary change: schools and workplaces. *J. Nutr.* 135:910-12
42. French S, Jeffery R, Story M, Breitlow K, Baxter J, et al. 2001. Pricing and promotion effects on low-fat vending snack purchases: the CHIPS Study. *Am. J. Public Health* 91:112-17
43. French S, Lin B, Guthrie J. 2003. National trends in soft drink consumption among children and adolescents age 6 to 17 years: prevalence, amounts, and sources, 1977-1978 to 1994-1998. *J. Am. Diet. Assoc.* 103:1326-31
44. French S, Story M, Jeffery R, Snyder P, Eisenberg M, et al. 1997. Pricing strategy to promote fruit and vegetable purchase in high school cafeterias. *J. Am. Diet. Assoc.* 97:1008-10
45. French SA, Story M, Jeffery RW. 2001. Environmental influences on eating and physical activity. *Annu. Rev. Public Health* 22:309-35
46. Gallo A. 1999. *America's Eating Habits: Changes and Consequences*, Washington, DC: Econ. Res. Serv., USDA
47. Glanz K, Hoelscher D. 2004. Increasing fruit and vegetable intake by changing environments, policy and pricing: restaurant-based research, strategies, and recommendations. *Prev. Med.* 39:S88-93
48. Glanz K, Lankenau B, Foerster S, Temple S, Mullis R, Schmid T. 1995. Environmental and policy approaches to cardiovascular disease prevention through nutrition: opportunities for state and local action. *Health Educ. Q.* 22:512-27
49. Glanz K, Yaroch A. 2004. Strategies for increasing fruit and vegetable intake in grocery stores and communities: policy, pricing, and environmental change. *Prev. Med.* 39:S75-80
50. Green K, Steer S, Maluk R, Mahaffey S, Muhajarine N. 1993. Evaluation of the Heart Smart Restaurant Program in

- Saskatoon and Regina, Saskatchewan. *Can. J. Public Health* 84:399–402
51. Gruber J. 2000. *Youth Smoking in the U.S.: Prices and Policies. Work. Pap. #7506*. Cambridge, MA: Natl. Bur. Econ. Res.
 52. Haire-Joshu D, Nanney M. 2002. Prevention of overweight and obesity in children: influences on the food environment. *Diabetes Educ.* 28:415–23
 53. Haire-Joshu D, Nanney MS. 2003. Behavioral interventions for cancer prevention: dietary intake and physical activity. In *Evidence-Based Interventions In Oncology*, ed. B Givens, B Givens, pp. 17–62. St. Louis, MO: Springer
 54. Hannan P, French SA, Story M, Fulkeron JA. 2002. A pricing strategy to promote sales of lower fat foods in high school cafeterias: acceptability and sensitivity analysis. *Am. J. Health Promot.* 17:1–6
 55. Harris K, Paine-Andrews A, Kichter K. 1997. Reducing elementary school children's risks for chronic diseases through school lunch modifications, nutrition education, and physical activity interventions. *J. Nutr. Educ.* 29:196–202
 56. Hearn M, Baranowski J, Doyle C, Smith M, Lin L, Resnicow K. 1998. Environmental influences on dietary behavior among children: availability and accessibility of fruits and vegetables enable consumption. *J. Health Educ.* 29:26–32
 57. Heath GW, Brownson RC, Kruger J, Miles R, Powell KE, et al. 2005. The effectiveness of environmental and policy interventions to increase physical activity: a systematic review. *Am. J. Prev. Med.* In press
 58. Hoelscher DM, Feldman HA, Johnson CC, Lytle LA, Osganian SK, et al. 2004. School-based health education programs can be maintained over time: results from the CATCH Institutionalization study. *Prev. Med.* 38:594–606
 59. Hoerr S, Loudon V. 1993. Can nutrition information increase sales of healthful vended snacks? *J. Sch. Health* 63:386–90
 60. Hopkins DP, Briss PA, Ricard CJ, Husten CG, Carande-Kulis VG, et al. 2001. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am. J. Prev. Med.* 20:16–66
 61. Humpel N, Owen N, Leslie E. 2002. Environmental factors associated with adults' participation in physical activity. A review. *Am. J. Prev. Med.* 22:188–99
 62. Hyland A, Higbee C, Bauer JE, Giovino GA, Cummings KM. 2004. Cigarette purchasing behaviors when prices are high. *J. Public Health Manag. Pract.* 10:497–500
 63. Israel BA, Schulz AJ, Parker EA, Becker AB. 1998. Review of community-based research: assessing partnership approaches to improve public health. *Annu. Rev. Public Health* 19:173–202
 64. James J, Thomas P, Cavan D, Kerr D. 2004. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomized controlled trial. *BMJ* 328:1237–50
 65. Jeffery R, Utter J. 2003. The changing environment and population obesity in the United States. *Obes. Res.* 11:12S–22S
 66. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, et al. 2002. The effectiveness of interventions to increase physical activity. A systematic review. *Am. J. Prev. Med.* 22:73–107
 67. Kaiser HJ. 2004. *The Role of Media in Childhood Obesity*. Menlo Park, CA: Henry J. Kaiser Found.
 68. Kerner J, Guirguis-Blake J, Hennessy K, Brounstein P, Vinson C, et al. 2006. Translating research into improved outcomes in comprehensive cancer control. *Cancer Causes Control*. In press
 69. Kerr J, Eves FF, Carroll D. 2001. The influence of poster prompts on stair use: The effects of setting, poster size and content. *Br. J. Health Psychol.* 6:397–405
 70. King AC, Jeffery RW, Fridinger F, Dusenbury L, Provence S, et al. 1995. Environmental and policy approaches to cardiovascular disease prevention through

- physical activity: issues and opportunities. *Health Educ. Q.* 22:499–511
71. King C 3rd, Siegel M. 2001. The Master Settlement Agreement with the tobacco industry and cigarette advertising in magazines. *N. Engl. J. Med.* 345:504–11
72. Krugman DM, Fox RJ, Fischer PM. 1999. Do cigarette warnings warn? Understanding what it will take to develop more effective warnings. *J. Health Commun.* 4:95–104
73. Kuchler F, Tegene A, Harris J. 2004. Taxing snack foods: what to expect for diet and tax revenues. *Agric. Inf. Bul.*:1–12
74. Kuchler F, Tegene A, Harris J. 2005. Taxing snack foods: manipulating diet quality or financing information programs? *Rev. Agric. Econ.* 27:4–20
75. Last JM, ed. 2001. *A Dictionary of Epidemiology*. New York: Oxford Univ. Press. 196 pp.
76. Ledikwe J, Ello-Martin J, Rolls B. 2005. Portion sizes and the obesity epidemic. *J. Nutr.* 135(4):905–9
77. Levy DT, Chaloupka F, Gitchell J. 2004. The effects of tobacco control policies on smoking rates: a tobacco control scorecard. *J. Public Health Manag. Pract.* 10:338–53
78. Lewit E, Coate D, Grossman M. 1981. The effects of government regulation on teenage smoking. *J. Law Econ.* 24:545–69
79. Lin B, Frazao E, Guthrie J. 1999. *Away from Home Foods Increasingly Important to Quality of American Diet. Rep. No. 749*. Washington, DC: ARS, USDA
80. Luke D. 2005. Getting the big picture in community science: methods that capture context. *Am. J. Community Psychol.* 35:185–200
81. Luke DA, Stamatakis KA, Brownson RC. 2000. State youth access tobacco control policies and youth smoking behavior in the United States. *Am. J. Prev. Med.* 19:180–87
82. Lytle L, Murray D, Perry C, Story M, Birnbaum A, et al. 2004. School-based approaches to affect adolescents' diets: results from the TEENS study. *Health Educ. Behav.* 31:270–87
83. Martens D. 2002. Graphic tobacco warnings having desired effect. *CMAJ* 166:1453
84. Martinez RM, McHugh M, Kliman R, Roschwalb S. 1999. *Community Preventive Services in Ten Health Departments and Their Receptivity to Evidence-Based Guidelines*. Washington, DC: Math. Policy Res.
85. Matson-Koffman DM, Brownstein JN, Neiner JA, Greaney ML. 2005. A site-specific literature review of policy and environmental interventions that promote physical activity and nutrition for cardiovascular health: What works? *Am. J. Health. Promot.* 19:167–93
86. McKenna MT, Taylor WR, Marks JS, Koplan JP. 1998. Current issues and challenges in chronic disease control. In *Chronic Disease Epidemiology and Control*, ed. RC Brownson, PL Remington, JR Davis, pp. 1–26. Washington, DC: Am. Public Health Assoc.
87. Meyer M, Miller E. 1984. *Urban Transportation Planning: A Decision Oriented Approach*. New York: McGraw-Hill
88. Mitra A. 2001. Effects of physical attributes on the wages of males and females. *Appl. Econ. Lett.* 8:731–35
89. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. 2004. Actual causes of death in the United States, 2000. *JAMA* 291:1238–45
90. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. 2005. Correction: actual causes of death in the United States, 2000. *JAMA* 293:293–94
91. Mudarri D. 1994. *The Costs and Benefits of Smoking Restrictions: an Assessment of the Smoke-Free Environment Act of 1993 (H.R. 3434)*. Rep. Gov. Doc. 1994; H.R. 3434, Washington, DC: US GPO
92. Muller PO. 1995. Transportation and urban form: stages in the spatial evolution

- of the American metropolis. In *The Geography of Urban Transportation*, ed. S Hanson, pp. 26–52. New York: Guilford
93. Natl. Cancer Inst. 2000. *Health Effects of Exposure to Environmental Tobacco Smoke. Smoking and Tobacco Control Monographs 10*. Bethesda, MD: NCI
 94. Natl. Cent. Health Stat. 1999. *Healthy People 2000 Review 1998–1999. Rep. DHHS Publ No (PHS) 99–1256*, Hyattsville, MD: US DHHS
 95. Nicklas N, Johnson C, Myers L. 1998. Outcomes of a high school program to increase fruit and vegetable consumption: gimme 5—a fresh nutrition concept for students. *J. Sch. Health* 68:248–53
 96. Nielsen S, Popkin B. 2003. Patterns and trends in food portion sizes, 1977–1998. *J. Am. Med. Assoc.* 289:450–53
 97. Nike. 2005. *NikeGO*. <http://www.nike.com/nikebiz/nikego/index.html>
 98. Owen N, Humpel N, Leslie E, Bauman A, Sallis JF. 2004. Understanding environmental influences on walking: review and research agenda. *Am. J. Prev. Med.* 27:67–76
 99. Perlmutter CA, Canter DD, Gregoire MB. 1997. Profitability and acceptability of fat- and sodium-modified hot entrees in a worksite cafeteria. *J. Am. Diet. Assoc.* 97:391–95
 100. Perry C, Bishop D, Taylor G. 1998. Changing fruit and vegetable consumption among children: the 5-a-Day Power Plus Program in St. Paul, Minnesota. *Am. J. Public Health* 88:603–9
 101. Pratt M, Macera CA, Sallis JF, O'Donnell M, Frank LD. 2004. Economic interventions to promote physical activity: application of the SLOTH model. *Am. J. Prev. Med.* 27:136–45
 102. Pratt M, Macera CA, Wang G. 2000. Higher direct medical costs associated with physical inactivity. *Phys. Sportsmed.* 28:63–70
 103. Raynor H, Epstein L. 2001. Dietary variety, energy regulation and obesity. *Psychol. Bull.* 3:127–36
 104. Reger B, Wootan M, Booth-Butterfield S. 1999. Using mass media to promote healthy eating: a community based demonstration project. *Prev. Med.* 29:414–21
 105. Roussos ST, Fawcett SB. 2000. A review of collaborative partnerships as a strategy for improving community health. *Annu. Rev. Public Health* 21:369–402
 106. Saelens BE, Sallis JF, Frank LD. 2002. Environmental correlates of walking and cycling: Findings from the transportation, urban design and planning literatures. *Ann. Behav. Med.* 25:80–91
 107. Saffer H, Chaloupka F. 2000. The effect of tobacco advertising bans on tobacco consumption. *J. Health Econ.* 19:1117–37
 108. Sallis J, Cervero R, Ascher W, Henderson K, Kraft M, Kerr J. 2006. An ecological approach to creating active living communities. *Annu. Rev. Public Health* 27:297–322
 109. Sallis JF, Bauman A, Pratt M. 1998. Environmental and policy interventions to promote physical activity. *Am. J. Prev. Med.* 15:379–97
 110. Schilling J, Linton LS. 2005. The public health roots of zoning: in search of active living's legal genealogy. *Am. J. Prev. Med.* 28:96–104
 111. Schmid TL, Pratt M, Howze E. 1995. Policy as intervention: environmental and policy approaches to the prevention of cardiovascular disease. *Am. J. Public Health* 85:1207–11
 112. Scollo M, Youniss S, Wakefield M, Freeman J, Icasiano F. 2003. Impact of tobacco tax reforms on tobacco prices and tobacco use in Australia. *Tob. Control* 12(Suppl. 2):ii59–66
 113. Shadish W, Cook T, Campbell D. 2002. *Experimental and Quasi-Experimental Designs for Generalized Causal Inference*. Boston, MA: Houghton Mifflin
 114. Siegel M. 2002. The effectiveness of state-level tobacco control interventions: a review of program implementation and behavioral outcomes. *Annu. Rev. Public Health* 23:45–71

115. Sorensen G, Stoddard A, Peterson K, Cohen N, Hunt M. 1999. Increasing fruit and vegetable consumption through work-sites and families in the Treatwell 5-a-day study. *Am. J. Public Health* 89:54–60
116. Spasoff RA. 1999. *Epidemiologic Methods for Health Policy*. New York: Oxford Univ. Press
117. Stables G, Subar A, Patterson B, Dodd K, Heimendinger J, et al. 2002. Changes in fruit and vegetable consumption and awareness among US adults: Results of the 1991 and 1997 5-a-day for better health program surveys. *J. Am. Diet. Assoc.* 102:809–17
118. Stahl T, Rutten A, Nutbeam D, Kannas L. 2002. The importance of policy orientation and environment on physical activity participation—a comparative analysis between Eastern Germany, Western Germany and Finland. *Health Promot. Int.* 17:235–46
119. Story M, Neumark-Sztainer D, French S. 2002. Individual and environmental influences on adolescent eating behaviors. *J. Am. Diet. Assoc.* 102:S40–51
120. Subramanian SV, Belli P, Kawachi I. 2002. The macroeconomic determinants of health. *Annu. Rev. Public Health* 23:287–302
121. Tauras J, Chaloupka F. 1999. *Price, Clean Indoor Air Laws, and Cigarette Smoking: Evidence from Longitudinal Data for Young Adults*. Work. Pap. #6937. Cambridge, MA: Natl. Bur. Econ. Res.
122. Thacker SB, Koplan JP, Taylor WR, Hinman AR, Katz MF, Roper WL. 1994. Assessing prevention effectiveness using data to drive program decisions. *Public Health Rep.* 109:187–94
123. Tillotson J. 2003. Pandemic obesity: unintended policy consequences. *Nutr. Today* 38:116–19
124. Tillotson J. 2004. America's obesity: conflicting public policies, industrial economic development, and unintended human consequences. *Annu. Rev. Nutr.* 24: 617–43
125. Trevino R, Pugh J, Hernandez A. 1998. Bienstar: a diabetes risk-factor prevention program. *J. Sch. Health* 68:62–67
126. Truman BI, Smith-Akin CK, Hinman AR, et al. 2000. Developing the guide to community preventive services—overview and rationale. *Am. J. Prev. Med.* 18:18–26
127. Turnock BJ. 2001. *Public Health: What it is and How it Works*. Gaithersburg, MD: Aspen Publ. 354 pp.
128. US Dep. Health Hum. Serv. 1986. *The Health Consequences of Involuntary Smoking: A Report of the Surgeon General*. Rep. 97–8398, Washington, DC: US DHHS
129. US Dep. Health Hum. Serv. 1994. *Preventing Tobacco Use Among Young People: A Report of the Surgeon General*. Atlanta, GA: Cent. Dis. Control Prev.
130. van Wechem S, Brug J, van Assema P, Kistemaker C, Riedstra M, Lowik M. 1998. Fat Watch: a nationwide campaign in The Netherlands to reduce fat intake—effect evaluation. *Nutr. Health* 12:119–30
131. Veugelers P, Fitzgerald A. 2005. Effectiveness of school programs in preventing childhood obesity: a multilevel comparison. *Am. J. Public Health* 95:431–35
132. Wakefield M, Giovino G. 2003. Teen penalties for tobacco possession, use, and purchase: evidence and issues. *Tob. Control* 12 (Suppl. 1):i6–13
133. Wang G, Macera CA, Scudder-Soucie B, Schmid T, Pratt M, et al. 2004. Cost analysis of the built environment: the case of bike and pedestrian trails in Lincoln, Neb. *Am. J. Public Health* 94:549–53
134. Wansink B. 2004. Environmental factors that increase the food intake and consumption volume of unknowing consumers. *Annu. Rev. Nutr.* 24:455–79
135. Wasserman J, Manning WG, Newhouse JP, Winkler JD. 1991. The effects of excise taxes and regulations on cigarette smoking. *J. Health Econ.* 10:43–64
136. Webb K, Hawe P, Noort M. 2001.

- Collaborative intersectoral approaches to nutrition in a community on the urban fringe. *Health Educ. Behav.* 28:306–19
137. Wechsler H, Brener N, Kuester S, Miller C. 2001. Food service and foods and beverages available at school: results from the School Health Policies and Programs Study 2000. *J. Sch. Health* 71:313–24
138. Wiggers J, Considine R, Hazell T, Haile M, Rees M, Daly J. 2001. Increasing the practice of health promotion initiatives by licensed premises. *Health Educ. Behav.* 28:331–40
139. Wilkinson-Enns C, Goldman J, Cook A. 1997. Trends in food and nutrient intakes by adults: NFCS 1977–78, CSFII 1989–91, and CSFII 1994–95. *Fam. Econ. Nutr. Rev.* 10:2–15
140. World Health Organ. 2005. *Global Strategy on Diet, Physical Activity and Health*. Geneva: WHO
141. Yach D, Hawkes C, Gould CL, Hofman KJ. 2004. The global burden of chronic diseases: overcoming impediments to prevention and control. *JAMA* 291:2616–22
142. Yach D, McKee M, Lopez AD, Novotny T. 2005. Improving diet and physical activity: 12 lessons from controlling tobacco smoking. *BMJ* 330:898–900
143. Young L, Nestle M. 2002. The contribution of expanding portion sizes to the US obesity epidemic. *Am. J. Public Health* 92:246–49
144. Young L, Nestle M. 2003. Expanding portion sizes in the US marketplace: Implication for nutrition counseling. *J. Am. Diet. Assoc.* 103:231–34
145. Zaza S, Briss PA, Harris KW, eds. 2005. *The Guide to Community Preventive Services: What Works to Promote Health?* New York: Oxford Univ. Press



CONTENTS

EPIDEMIOLOGY AND BIostatISTICS

- Effective Recruitment and Retention of Minority Research Participants,
Antronette K. Yancey, Alexander N. Ortega, and Shiriki K. Kumanyika 1
- Measuring Population Health: A Review of Indicators, *Vera Etches,
John Frank, Erica Di Ruggiero, and Doug Manuel* 29
- On Time Series Analysis of Public Health and Biomedical Data,
Scott L. Zeger, Rafael Irizarry, and Roger D. Peng 57
- The Promise and Pitfalls of Systematic Reviews, *Patricia Dolan Mullen
and Gilbert Ramírez* 81
- Hypertension: Trends in Prevalence, Incidence, and Control, *Ihab Hajjar,
Jane Morley Kotchen, and Theodore A. Kotchen* 465

ENVIRONMENTAL AND OCCUPATIONAL HEALTH

- Environmental Justice: Human Health and Environmental Inequalities,
Robert J. Brulle and David N. Pellow 103
- Speed, Road Injury, and Public Health, *Elihu D. Richter, Tamar Berman,
Lee Friedman, and Gerald Ben-David* 125
- The Big Bang? An Eventful Year in Workers' Compensation,
Tee L. Guidotti 153
- Shaping the Context of Health: A Review of Environmental and Policy
Approaches in the Prevention of Chronic Diseases, *Ross C. Brownson,
Debra Haire-Joshu, and Douglas A. Luke* 341

PUBLIC HEALTH PRACTICE

- Health Disparities and Health Equity: Concepts and Measurement,
Paula Braveman 167
- The Politics of Public Health Policy, *Thomas R. Oliver* 195
- Vaccine Shortages: History, Impact, and Prospects for the Future,
*Alan R. Hinman, Walter A. Orenstein, Jeanne M. Santoli,
Lance E. Rodewald, and Stephen L. Cochi* 235
- What Works, and What Remains to Be Done, in HIV Prevention in the
United States, *David R. Holtgrave and James W. Curran* 261

SOCIAL ENVIRONMENT AND BEHAVIOR

- A Public Health Success: Understanding Policy Changes Related to Teen Sexual Activity and Pregnancy, *Claire D. Brindis* 277
- An Ecological Approach to Creating Active Living Communities, *James F. Sallis, Robert B. Cervero, William Ascher, Karla A. Henderson, M. Katherine Kraft, and Jacqueline Kerr* 297
- Process Evaluation for Community Participation, *Frances Dunn Butterfoss* 323
- Shaping the Context of Health: A Review of Environmental and Policy Approaches in the Prevention of Chronic Diseases, *Ross C. Brownson, Debra Haire-Joshu, and Douglas A. Luke* 341
- Stress, Fatigue, Health, and Risk of Road Traffic Accidents Among Professional Drivers: The Contribution of Physical Inactivity, *Adrian H. Taylor and Lisa Dorn* 371
- The Role of Media Violence in Violent Behavior, *L. Rowell Huesmann and Laramie D. Taylor* 393

HEALTH SERVICES

- Aid to People with Disabilities: Medicaid's Growing Role, *Alicia L. Carbaugh, Risa Elias, and Diane Rowland* 417
- For-Profit Conversion of Blue Cross Plans: Public Benefit or Public Harm? *Mark A. Hall and Christopher J. Conover* 443
- Hypertension: Trends in Prevalence, Incidence, and Control, *Ihab Hajjar, Jane Morley Kotchen, and Theodore A. Kotchen* 465
- Preventive Care for Children in the United States: Quality and Barriers, *Paul J. Chung, Tim C. Lee, Janina L. Morrison, and Mark A. Schuster* 491
- Public Reporting of Provider Performance: Can Its Impact Be Made Greater? *David L. Robinowitz and R. Adams Dudley* 517
- Health Disparities and Health Equity: Concepts and Measurement, *Paula Braveman* 167

INDEXES

- Subject Index 537
- Cumulative Index of Contributing Authors, Volumes 18–27 565
- Cumulative Index of Chapter Titles, Volumes 18–27 570

ERRATA

An online log of corrections to *Annual Review of Public Health* chapters may be found at <http://publhealth.annualreviews.org/>